

Does Job Stress Affect Injury Due to Labor Accident in Japanese Male and Female Blue-Collar Workers?

Katsuyuki MURATA^{1*}, Norito KAWAKAMI² and Noriko AMARI¹

¹Department of Hygiene and Public Health, Teikyo University School of Medicine, 2–11–1 Kaga, Itabashi-ku, Tokyo 173-8605, Japan

²Department of Public Health, Gifu University School of Medicine, 40 Tsukasa-machi, Gifu 500-8705, Japan

Received September 20, 1999 and accepted January 31, 2000

Abstract: To clarify whether job stressors affecting injury due to labor accidents differ between Japanese male and female blue-collar workers, the Job Content Questionnaire (JCQ), assessing dimensions of job stressors based on the demand-control-support model, was applied to 139 blue-collar workers in a manufacturing factory. Of them, 24 male and 15 female workers suffered from injuries at work. In the female workers with the experience of work injury, the job demand score and job strain index (i.e., the ratio of job demand to job control) of the JCQ were significantly higher and the score of coworker support was significantly lower, than those in the female workers without the experience. High job demand (or, high job strain and low coworker support) was significantly related to work injury in all the female workers. Between the male workers with and without work injury, however, there was no significant difference in any job stressors. This pilot study suggests that high job strain (specifically, high job demand), as well as low coworker support, are important factors affecting work injury in Japanese female blue-collar workers. Further research with a large number of male blue-collar workers will be required to seek other factors that may be associated with work injury.

Key words: Work injury, Job stress, Job content questionnaire, Blue-collar workers, Gender

Introduction

Little information has been developed regarding whether the relation of job stress to injury due to labor accident differs between male and female workers, although the effects of job stress are well-known to vary by sex^{1,2}. Certainly, psychosocial factors at work appear to affect work-related injuries including musculoskeletal diseases³, because the involvement of the human element in accident and injury causation is ubiquitous⁴ and the majority of work injuries have some contributory cause related to the interaction between the worker and the work system⁵. Moreover, there is evidence that occupational factors that cause psychological strain can affect physical and mental health such as cardiovascular disease^{6,7}, epicondylitis⁸, psychiatric disorders^{9–11}, and alcohol dependence¹²; whereas,

the findings are not always consistent^{1,13}. Recently, Wilkins and Beaudet have reported a sex difference in the link between work injury and job strain, with use of the data obtained from the Canada's 1994/95 National Population Health Survey¹⁴. The definition of "work injury" may, however, have varied across individuals. For worksite health promotion, the sex difference in such associations should be verified in countries with different cultures.

For assessing job stressors, two well-established questionnaires, i.e., the Job Content Questionnaire (JCQ)¹⁵ and Generic Job Stress Questionnaire (GJSQ)^{16,17}, are commonly used in the world. These two also have relatively high reliability and validity¹³. In this article, a cross-sectional study was conducted to clarify whether job stressors, based on the demand-control-support model¹⁵, affecting injury due to labor accident differ between Japanese male and female blue-collar workers.

*To whom correspondence should be addressed.

Methods

Subjects

In March 1999, a self-rating questionnaire was distributed to all of 201 full-time employees of a small manufacturing factory in Japan (approximately 80 part-time and seasonal laborers were excluded because their working duration was relatively short and their occupational histories were not fully followed up). The response rate to this questionnaire was 83.6% (of 33 non-respondents, eleven were white-collar workers, and three of the remaining 22 blue-collar workers had the experience of work injury). The study cohort, accordingly, consisted of 92 male and 76 female workers, aged 19–59 (mean 40) years. In them, two men with diabetes mellitus and ten men with hypertension were included. This factory manufactured many different kinds of aerosol products seasonally, and the work system and work environment had been basically unchanged despite some minor improvements, at least for ten years. Seventy-six men and 63 women had worked in divisions of solvent-blending, aerosol-injection, bulb-assembling, packing, carriage of goods, and machinery maintenance (i.e., blue-collar workers). Twenty-nine workers were engaged in administrative/clerical work, or research (i.e., white-collar workers). None of the 168 workers had changed their occupation between blue-collar and white-collar, but a few blue-collar workers shifted among production divisions when they could not accommodate themselves to the work absolutely.

Methods

The JCQ, developed by Karasek¹⁵⁾, has been translated into Japanese and its factor-based and construct validity and internal consistency reliability have been tested in several samples of Japanese workers¹⁸⁾. The questionnaire included 22 questions selected from the full recommended version (45 items) of the JCQ. These items constituted a minimum set of questions for four major scales of the JCQ: (1) job demand (five items), (2) job control or decision latitude (nine items), (3) supervisor support (four items), and (4) coworker support (four items). Each of the items had a 4-point response ranging from 1 (strongly disagree) to 4 (strongly agree). According to the Scoring Manual of the JCQ¹⁵⁾, a sum of weighted item scores was used as a scale score. Also, job strain index was defined as the score ratio of job demand to job control.

All injuries due to labor accidents in the factory (except traffic accidents in commute) had been recorded by an occupational health nurse since April, 1989. As shown in Table 1, this study included not only slight (so-called

Table 1. Labor accidents in a manufacturing factory during the period of April, 1989 to March, 1999^a

	Male workers	Female workers
Part of body injured		
Fingers	23	11
Legs/feet	4	5
Trunk	1	7
Head	6	1
Eyes	5	2
Arms	4	2
Hands	3	3
General	1	2
Injuries		
Contused wound	18	10
Contusion	9	14
Incised wound	10	1
Bone fracture	4	4
Injuries	4	2
Burn	1	1
Dermatitis	1	0
Low back pain	0	1

^a Accident-based number. Resignees were included in these figures, but part-time and seasonal laborers were excluded.

“*Akachin* (iodine tincture)”) injuries such as contusion and incised wound, but also moderate and severe injuries which confined the worker to bed for more than half of the daylight hours on the date of the accident or following days (e.g., bone fracture). On the accident-basis, the proportion of the “*Akachin*” injury to a total of labor accidents for the period of April, 1989 to March, 1999, was 27% both in males and females including resignees and accident-prone workers who experienced such accidents twice or over. Of the 139 blue-collar workers, 24 males and 15 females suffered from injuries in the workplace; seven males and two females were accident-prone workers (29% and 13%, respectively).

Data analysis

The comparison of job stress scores between the workers with and without the experience of work injury was made by the Student's *t* test and analysis of covariance. Sex difference was tested by the analysis of variance (ANOVA) with repeated measurements (SS model of type II). Also, the multiple logistic regression analysis, including age and working duration as confounders (i.e., independent variables), was used to examine which of job demand, job control, and supervisor and coworker support was closely associated with the presence/absence of work injury. All the analyses were performed using the Statistical Package for the Biosciences¹⁹⁾.

Table 2. Job stress scores of the Job Content Questionnaire in blue-collar and white-collar workers in a manufacturing factory (mean \pm SD)

	Blue-collar workers	White-collar workers	Probability (P) by t test
Males (Number):	(76)	(16)	
Job demand	33.0 \pm 5.1	34.2 \pm 3.2	0.237 (0.471)*
Job control	63.6 \pm 9.6	71.6 \pm 8.7	0.003 (0.002)*
Supervisor support	11.2 \pm 2.2	11.9 \pm 2.7	0.308 (0.343)*
Coworker support	11.4 \pm 1.7	12.2 \pm 1.8	0.088 (0.111)*
Job strain index	0.531 \pm 0.120	0.483 \pm 0.062	0.026 (0.096)*
Age (years)	37.5 \pm 11.2	35.1 \pm 12.6	0.456
Working duration (years)	13.9 \pm 9.8	10.2 \pm 10.6	0.171 (0.249)*
Females (Number):	(63)	(13)	
Job demand	32.9 \pm 5.4	30.5 \pm 5.6	0.147 (0.497)*
Job control	53.5 \pm 10.3	65.8 \pm 8.4	<0.001 (0.002)*
Supervisor support	10.5 \pm 2.8	11.2 \pm 3.1	0.482 (0.573)*
Coworker support	11.3 \pm 2.0	12.0 \pm 2.2	0.299 (0.812)*
Job strain index	0.639 \pm 0.169	0.465 \pm 0.079	<0.001 (0.018)*
Age (years)	46.1 \pm 12.4	28.8 \pm 12.1	<0.001
Working duration (years)	9.5 \pm 7.3	6.6 \pm 5.7	0.180 (0.599)*

*Probability by the analysis of covariance used to control for age.

Results

The job stress scores of the JCQ in white-collar and blue-collar workers are shown in Table 2. In both male and female workers, the score of job control was significantly lower in the blue-collar workers than in the white-collar workers (ANOVA, $P < 0.001$). On the other hand, the job strain index was significantly higher in the blue-collar workers (ANOVA, $P < 0.001$). Significant sex differences were also found in job control and job strain index (ANOVA, $P < 0.001$).

In the female blue-collar workers with the experience of work injury, the job demand score and job strain index were significantly higher than those in the female workers without the experience (Table 3). Similarly, the coworker support score was significantly lower in the female blue-collar workers. Although all the scores of job stressors in the female blue-collar workers who suffered from injuries were comparable between two 5-year periods (i.e., of April, 1989 to March, 1994, and of April, 1994 to March, 1999), these subgroups had higher scores of job demand as compared to the women without the experience of work injury (Table 3). On the other hand, there was no significant difference between the male blue-collar workers with and without the experience of work injury (Table 3). Among three subgroups of the male blue-collar workers, i.e., workers without an accident, those with one accident and those with two or more accidents, no significant difference was found in any job stressors (one-way analysis of variance, $P > 0.05$).

High job demand was significantly related to work injury in the female blue-collar workers, but any job stressors were not significantly related in the male blue-collar workers (Table 4). When the job strain index, instead of job demand and job control scales, was used as one of independent variables of the multiple logistic regression analysis, job strain (estimated $\beta = 6.00$, standard error = 2.18, $P = 0.006$) and coworker support (estimated $\beta = -0.349$, standard error = 0.178, $P = 0.050$) were significantly related to work injury in the female blue-collar workers, but any stressors were not related in the males.

Discussion

In this study, the female blue-collar workers with the experience of work injury had high job demand and job strain, and low coworker support as compared to those without such an experience, although the job control score was comparable. In addition, high job demand (or, high job strain and low coworker support) was significantly associated with work injury in the female blue-collar workers. In Canadian female workers, high job strain measured as a ratio of psychological demands to decision latitude, and low coworker support were linked with work injury in the past 12 months¹⁴; which of the psychological demand and decision latitude contributed to high job strain has not been discussed in the paper. Workers with back injury in Baltimore, US, also had a close relation to high job strain²⁰. In a 5-

Table 3. Job stress scores in blue-collar workers with and without the experience of work injury (mean ± SD)

	Workers with experience of work injury			Workers without experience of work injury	Probability* (P)	
	Total	in 1989–94 ^e	in 1994–99 ^e		age-adjusted ^a	duration-adjusted ^b
Males (Number):	(24)	(11)	(13)	(52)		
Job demand	33.5 ± 5.5	33.9 ± 6.0	33.1 ± 6.0	32.8 ± 5.0	0.662	0.554
Job control	61.3 ± 8.3	60.5 ± 9.4	62.0 ± 7.6	64.6 ± 10.0	0.176	0.161
Supervisor support	10.8 ± 2.1	10.8 ± 2.3	10.7 ± 2.0	11.4 ± 2.3	0.211	0.202
Coworker support	11.1 ± 2.0	11.2 ± 2.2	11.1 ± 1.8	11.5 ± 1.6	0.382	0.430
Job strain index	0.560 ± 0.148	0.581 ± 0.166	0.543 ± 0.136	0.517 ± 0.104	0.163	0.126
Age (years)	36.7 ± 9.4	37.8 ± 7.4	35.8 ± 11.0	37.8 ± 12.0		0.687 ^d
Working duration (years)	14.6 ± 8.1	16.2 ± 6.7	13.2 ± 9.1	13.7 ± 10.5		0.703 ^d
Females (Number):	(15)	(6)	(9)	(48)		
Job demand	37.1 ± 4.1	37.8 ± 4.3 ^e	36.6 ± 4.2 ^e	31.6 ± 5.0	<0.001	<0.001
Job control	51.1 ± 11.9	53.0 ± 7.9	49.8 ± 14.2	54.3 ± 9.8	0.377	0.356
Supervisor support	9.8 ± 3.6	8.8 ± 3.4	10.4 ± 3.9	10.8 ± 2.5	0.292	0.290
Coworker support	10.3 ± 2.3	9.5 ± 2.4 ^e	10.9 ± 2.2	11.7 ± 1.8	0.046	0.025
Job strain index	0.753 ± 0.151	0.726 ± 0.131	0.771 ± 0.168 ^e	0.603 ± 0.159	0.005	0.003
Age (years)	50.3 ± 9.5	47.2 ± 11.8	52.4 ± 7.7	44.7 ± 13.0		0.129 ^d
Working duration (years)	11.0 ± 3.9	11.7 ± 4.6	10.6 ± 3.6	9.0 ± 8.0		0.208 ^d

*Probability by the analysis of covariance, between two groups of workers with and without experience of work injury, used to control for either age (^a) or working duration (^b). ^cThe latest year was adopted in case of the accident-prone worker; and, one way analysis of variance was made among workers with experience of work injuries in 1989–94, and in 1994–99, and those without the experience. ^dStudent’s t test (or Welch t test). ^eP<0.05 (v.s. workers without injury, Scheffe’s multiple comparison test).

Table 4. Relations of job demand, job control, supervisor and coworker support scores, age and working duration to the presence/absence of work injury in blue-collar workers: results of multiple logistic regression analysis

Independent variables	Estimated β	Standard error	Probability (P) by Wald test	Odds ratio
Males (N=76):				
Job demand	0.02627	0.05189	0.613	1.03
Job control	- 0.03992	0.02869	0.164	0.96
Supervisor support	- 0.15204	0.12937	0.240	0.86
Coworker support	- 0.01667	0.17332	0.923	0.98
Age	- 0.03723	0.03486	0.286	0.96
Working duration	0.04743	0.03994	0.235	1.05
Females (N=63):				
Job demand	0.26409	0.08331	0.002	1.30
Job control	- 0.05845	0.03650	0.109	0.94
Supervisor support	0.02261	0.12049	0.855	1.02
Coworker support	- 0.34609	0.18003	0.055	0.71
Age	0.01507	0.04177	0.718	1.02
Working duration	0.02747	0.05987	0.646	1.03

year prospective study of American urban transit operators, spinal injury was significantly predicted by psychological job demands, job dissatisfaction and the frequency of job problems²¹). Thus, the outcome of this pilot study suggests that high job strain (especially, high job demand) and low

coworker support are important factors affecting work injury in Japanese female blue-collar workers, as well as in Canadian female workers.

By contrast, no significant difference in job stressors was observed between the male blue-collar workers with and

without the experience of work injury in the present study. In Canadian male workers, likewise, none of job strain, job insecurity, coworker support and supervisor support were significantly related to work injury in past 12 months, while the male blue-collar workers showed a significantly high odds ratio of 3.2 to work injury¹⁴⁾.

The above results suggest that psychosocial factors affecting work injury may differ between both sexes. One reason for this difference may have been attributable to the fact that many female workers in Japan serve as assistants to male workers¹⁸⁾. Actually, job contents in the female blue-collar workers of this study were not always the same as those in the male blue-collar workers. In general, men are far more likely than women to have high control over their work process at the task level, and psychological job demands do not differ markedly between both sexes¹⁾; thus, job strain has been thought to be higher in female workers than in male workers. Nevertheless, it was not low job control but high job demand that was prominent in the female blue-collar workers with the experience of work injury of the current study. Female blue-collar workers who have more complaints about the way the work is organized, including the work flow and pace, the hours that are worked and the communication flow within the workplace, may be likely to suffer from work injury. On the other hand, since the incidence of work injury was higher in the male than in the female workers (31.6% for men and 23.8% for women, respectively) and there were more accident-prone workers in the male blue-collar workers than in the female, attention should be directed toward other factors such as individual characteristics including the type A behavior and mental ill health³⁾, together with high physical demands¹⁴⁾ that could not be examined in this study.

The comparability of the present study was preserved because age and working duration were comparable between the workers with and without work injury. All the subjects worked in the same factory, and the analyses were made in men and women separately. Moreover, the results of the significant difference in the age-adjusted score of job control between white- and blue-collar workers (Table 2) are concordant with other studies with a large number of samples^{1, 18)}.

On the contrary, there may have been some limitations in this study: (1) One is selection bias. The blue-collar workers were divided into two groups with and without work injury on the basis of the record reported by one occupational health nurse. Although the record would cover the majority of injuries that occurred in the factory and not include any recall bias, some slight injuries such as scratch might not have

been declared by the propositus inasmuch as the Japanese workers generally tend to underreport. (2) The sample size, especially of workers with work injury, was too small. Moreover, the proportion of non-respondents in the current study was 16.4% (33 workers) and somewhat high as compared to other reports^{18, 22)}. (3) It is uncertain whether measures of chronic job stressors by the JCQ reflected the past 10-year psychological states fully, although job stress scores of the workers who suffered from work injury were comparable between the two periods of 1989–1994 and 1994–1999 (Table 3). (4) During the period of 1989–1999, about eight accident events per year occurred in the factory, mainly due to human errors (not skill-based errors but rule-based mistakes⁴⁾; for this reason, the company conducted safety education to strengthen the motivation to be safe, together with some minor improvements of the work system and environment. Further research with many blue-collar workers is necessary to confirm whether such factors, including the effects of safety education and workplace improvements, have strengthened or weakened the outcome of this study.

One goal of occupational health activities is to assess abnormal or excessive human stressors in the workplace, to eliminate the stress to improve performance and reliability, and to minimize error (e.g., accident and injury). In order to reduce work injury relevant to job stressors such as high job demand and low coworker support in vulnerable population (i.e., female workers), introduction of the participatory management system for workplace improvements²³⁾ and worksite stress reduction programs²⁴⁾ may be practical, and also enhance the functional quality of the workplace.

Acknowledgments

We wish to thank Prof. Eiji Yano and Dr. Koichi Nonaka, Department of Hygiene and Public Health, Teikyo University School of Medicine, and Dr. Takashi Haratani, National Institute of Industrial Health, for their valuable comments and suggestions.

References

- 1) Karasek R, Theorell T (1990) *Healthy work: Stress, productivity, and the reconstruction of working life*. Basic Books, New York.
- 2) Hellerstedt WL, Jeffery RW (1997) The association of job strain and health behaviours in men and women. *Int J Epidemiol* **26**, 575–83.
- 3) Bongers PM, de Winter CR, Kompier MAJ, Hildebrandt

- VH (1993) Psychosocial factors at work and musculoskeletal disease. *Scand J Work Environ Health* **19**, 297–312.
- 4) Feyer AM, Williamson A (1998) Occupational injury: Risk, prevention and intervention. Taylor & Francis, London.
 - 5) Budnick LD (1993) Human factors in occupational medicine. *J Occup Med* **35**, 587–97.
 - 6) Karasek R, Baker D, Marxer F, Ahlbom A, Theorell T (1981) Job decision latitude, job demands, and cardiovascular disease: a prospective study of Swedish men. *Am J Public Health* **71**, 694–705.
 - 7) Johnson JV, Hall EM (1988) Job strain, work place social support, and cardiovascular disease: a cross-sectional study of a random sample of the Swedish working population. *Am J Public Health* **78**, 1336–42.
 - 8) Ono Y, Nakamura R, Shimaoka M, Hiruta S, Hattori Y, Ichihara G, Kamijima M, Takeuchi Y (1998) Epicondylitis among cooks in nursery schools. *Occup Environ Med* **55**, 172–9.
 - 9) Iwata N, Okuyama Y, Kawakami Y, Saito K (1988) Psychiatric symptoms and related factors in a sample of Japanese workers. *Psychol Med* **18**, 659–63.
 - 10) Broadhead WE, Blazer DG, George LK, Tse CK (1990) Depression, disability days, and days lost from work in a prospective epidemiologic survey. *JAMA* **264**, 2524–8.
 - 11) Shigemi J, Mino Y, Tsuda T, Babazono A, Aoyama H (1997) The relationship between job stress and mental health at work. *Ind Health* **35**, 29–35.
 - 12) Kawakami N, Araki S, Haratani T, Hemmi T (1993) Relations of work stress to alcohol use and drinking problems in male and female employees of a computer factory in Japan. *Environ Res* **62**, 314–24.
 - 13) Kawakami N, Haratani T (1999) Epidemiology of job stress and health in Japan: review of current evidence and future direction. *Ind Health* **37**, 174–86.
 - 14) Wilkins K, Beaudet MP (1998) Work stress and health. *Health Rep* **10**, 47–62.
 - 15) Karasek R (1985) Job content questionnaire and user's guide. University of Massachusetts at Lowell, Lowell.
 - 16) Hurrell JJ Jr, McLaney MA (1988) Exposure to job stress—A new psychometric instrument. *Scand J Work Environ Health* **14** (suppl), 27–8.
 - 17) Haratani T, Kawakami N, Araki S (1993) Reliability and validity of the Japanese version of NIOSH Generic Job Stress Questionnaire. *Jpn J Ind Health* **35** (suppl), S214 (in Japanese).
 - 18) Kawakami N, Kobayashi F, Araki S, Haratani T, Furui H (1995) Assessment of job stress dimensions based on the job demands-control model of employees of telecommunication and electric power companies in Japan: reliability and validity of the Japanese version of the Job Content Questionnaire. *Int J Behav Med* **2**, 358–75.
 - 19) Winestem Institute of Community Medicine (1999) SPBS user's guide. Winestem Institute of Community Medicine, Toda (in Japanese).
 - 20) Myers AH, Baker SP, Li G, Smith GS, Wiker S, Liang KY, Johnson JV (1999) Back injury in municipal workers: a case-control study. *Am J Public Health* **89**, 1036–41.
 - 21) Krause N, Ragland DR, Fisher JM, Syme SL (1998) Psychosocial job factors, physical workload, and incidence of work-related spinal injury: a 5-year prospective study of urban transit operators. *Spine* **23**, 2507–16.
 - 22) Niemcryk SJ, Jenkins CD, Rose RM, Hurst MW (1987) The prospective impact of psychosocial variables on rates of illness and injury in professional employees. *J Occup Med* **29**, 645–52.
 - 23) Kogi K (1993) Workplace strategies for the control of work-related risks. *Environ Res* **63**, 88–94.
 - 24) Murphy LR (1984) Occupational stress management: a review and appraisal. *J Occup Psychol* **57**, 1–15.